



Real incidence of endoleaks after TEVAR

RG McWilliams
Liverpool, UK

Disclosure of Interest

Speaker name:

- I have the following potential conflicts of interest to report:
- Consulting
- Employment in industry
- Shareholder in a healthcare company
- Owner of a healthcare company
- Other(s)
- I do not have any potential conflict of interest

Key variables

- Pathology
- Experience
- Graft
- Surveillance schedule
- Completeness of surveillance
- Mode of surveillance
- Duration of follow up

Endoleaks after endovascular repair of thoracic aortic aneurysms

Shane S. Parmer et al

- 69 TEVAR
- Mean FU 17.3 months
- 20/69 (29%) endoleaks
- 8/20 Type 1 (12%)
- 7/20 Type 2 (12%)
- 4/20 Type 3 (6%)
- 1/20 more than one type

Endoleaks Following Endovascular Repair of Thoracic Aortic Aneurysm: Etiology and Outcomes

Jose P. Morales et al


- 200 TEVAR DTA (2001-2006)
- 75% atherosclerotic
- Mean follow-up 30 months
- 39 (19.5%) endoleaks
- Characterisation not always precise
- Three phase CT?

Endoleaks Following Endovascular Repair of Thoracic Aortic Aneurysm: Etiology and Outcomes

Jose P. Morales et al



J Endovasc Ther 2008;15:631–638



Endoleaks Following Endovascular Repair of Thoracic Aortic Aneurysm: Etiology and Outcomes

Jose P. Morales et al

- 39/200 developed endoleak
- 33 primary/8 secondary
- Type 1 11 primary 3 secondary (7%)
- Type 3 5 primary 2 secondary (3.5%)
- Type 2 13 primary 3 secondary (7%)
- Tx for most 1/3, mainly conservative for 2


Clinical significance of type II endoleaks after thoracic endovascular aortic repair

Bischoff MS et al

- 344 TEVAR (1997-2012)
- Type 2 in 30/344 – 8.7%
- Median FU 29.5 months
- Multiphase CT
- Reintervention 9/30 (30%) 5/9 LSA

Clinical significance of type II endoleaks after thoracic endovascular aortic repair

Bischoff MS et al



Mechanisms of Failure and Outcome of Secondary Surgical Interventions After Thoracic Endovascular Aortic Repair (TEVAR)

Julia Dumfarth et al

- 421 TEVAR (1996-2009)
- 21 secondary surgical interventions
- Type 1, retrograde type A, distal aneurysm and infection
- Median interval 24 months
- In-hospital mortality 19%
- “the majority of these events could have been avoided by a more strict indication.”

Thoracic Endovascular Aortic Repair (TEVAR) for the treatment of aortic diseases: a position statement....


Martin Grabenwoger et al


- **The majority of endoleaks can be avoided by careful selection** particularly with regard to important morphological details such as the length of the landing zone, use of multiple stents, length of overlapping segments as well as severe angulation and massive aortic calcification (porcelain aorta).


Incomplete Endograft Apposition to the Aortic Arch: Bird-Beak Configuration Increases Risk of Endoleak Formation after Thoracic Endovascular Aortic Repair

Takaya Ueda et al


Figure 1







Type 2



Type 2







Bridging stent dislocation




Investigation of endoleak




CEUS




IVUS



Rotational angiography




Rotational angiography




Rotational angiography




Rotational angiography




Rotational angiography





Rotational angiography



Possible 3b









Conclusions

- Endoleak rates high
- Type 1 ~ 10%, Type 2 ~ 10%, Type 3 ~ 5%, Overall 20-30%
- Classification can be imprecise
- US/CEUS not an option
- Avoidance – improved grafts, more overlap, patient selection